

- Your pet has been administered and/or prescribed an antibiotic to help prevent or treat an infection of his/her surgical site.
  - Please follow all label directions and give the medication with a meal.
  - Possible side effects include vomiting, diarrhea, anorexia or general gastrointestinal upset.
  - If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.
- 

- Your pet has been administered and/or prescribed an antibiotic to help prevent or treat an infection of his/her surgical site.
  - Please follow all label directions and give the medication with a meal.
  - Possible side effects include vomiting, diarrhea, anorexia or general gastrointestinal upset.
  - If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.
- 

- Your pet has been administered and/or prescribed an antibiotic to help prevent or treat an infection of his/her surgical site.
  - Please follow all label directions and give the medication with a meal.
  - Possible side effects include vomiting, diarrhea, anorexia or general gastrointestinal upset.
  - If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.
- 

- Your pet has been administered and/or prescribed an antibiotic to help prevent or treat an infection of his/her surgical site.
- Please follow all label directions and give the medication with a meal.
- Possible side effects include vomiting, diarrhea, anorexia or general gastrointestinal upset.
- If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.

- A doctor has noted that your animal has health issues of the mouth.
  - Problems in the mouth may cause your animal pain or lead to infection.
  - Please see your regular veterinarian for a thorough exam of your animal's mouth and any necessary treatment.
- 

- A doctor has noted that your animal has health issues of the mouth.
  - Problems in the mouth may cause your animal pain or lead to infection.
  - Please see your regular veterinarian for a thorough exam of your animal's mouth and any necessary treatment.
- 

- A doctor has noted that your animal has health issues of the mouth.
  - Problems in the mouth may cause your animal pain or lead to infection.
  - Please see your regular veterinarian for a thorough exam of your animal's mouth and any necessary treatment.
- 

- A doctor has noted that your animal has health issues of the mouth.
  - Problems in the mouth may cause your animal pain or lead to infection.
  - Please see your regular veterinarian for a thorough exam of your animal's mouth and any necessary treatment.
-

A doctor has noted that your animal has an ear issue. The cause can be ear mites (a tiny bug that feeds on the ear and is a common issue for kittens/young cats), allergies, yeast, bacteria, or foreign bodies. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable.

---

A doctor has noted that your animal has an ear issue. The cause can be ear mites (a tiny bug that feeds on the ear and is a common issue for kittens/young cats), allergies, yeast, bacteria, or foreign bodies. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable.

---

A doctor has noted that your animal has an ear issue. The cause can be ear mites (a tiny bug that feeds on the ear and is a common issue for kittens/young cats), allergies, yeast, bacteria, or foreign bodies. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable.

---

A doctor has noted that your animal has an ear issue. The cause can be ear mites (a tiny bug that feeds on the ear and is a common issue for kittens/young cats), allergies, yeast, bacteria, or foreign bodies. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable.

---

E-collars help prevent your pet from self-traumatizing and/or infecting their surgery site. They are bulky – and some dogs are somewhat distressed when initially wearing them – however, this is usually very temporary, and most pets quickly learn to tolerate them.

Your pet should have no trouble eating or drinking while wearing an E-collar, but if you observe them having difficulty doing so, you may remove it temporarily – you must, however, provide 100% supervision during this time.

Your dog should not be allowed to lick or otherwise rub at the surgery site, and must wear the E-collar for \_\_\_\_\_ days.

---

E-collars help prevent your pet from self-traumatizing and/or infecting their surgery site. They are bulky – and some dogs are somewhat distressed when initially wearing them – however, this is usually very temporary, and most pets quickly learn to tolerate them.

Your pet should have no trouble eating or drinking while wearing an E-collar, but if you observe them having difficulty doing so, you may remove it temporarily – you must, however, provide 100% supervision during this time.

Your dog should not be allowed to lick or otherwise rub at the surgery site, and must wear the E-collar for \_\_\_\_\_ days.

---

E-collars help prevent your pet from self-traumatizing and/or infecting their surgery site. They are bulky – and some dogs are somewhat distressed when initially wearing them – however, this is usually very temporary, and most pets quickly learn to tolerate them.

Your pet should have no trouble eating or drinking while wearing an E-collar, but if you observe them having difficulty doing so, you may remove it temporarily – you must, however, provide 100% supervision during this time.

Your dog should not be allowed to lick or otherwise rub at the surgery site, and must wear the E-collar for \_\_\_\_\_ days.

---

Feline Leukemia Virus (FeLV) is a viral infection in cats. There is no effective treatment to eliminate the infection. FeLV is spread from cat to cat through saliva, such as friendly grooming behavior or bite wounds, and is also spread from an infected mother cat to her kittens. Spaying eliminates this risk. If your cat tested positive, please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

Feline Leukemia Virus (FeLV) is a viral infection in cats. There is no effective treatment to eliminate the infection. FeLV is spread from cat to cat through saliva, such as friendly grooming behavior or bite wounds, and is also spread from an infected mother cat to her kittens. Spaying eliminates this risk. If your cat tested positive, please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

Feline Leukemia Virus (FeLV) is a viral infection in cats. There is no effective treatment to eliminate the infection. FeLV is spread from cat to cat through saliva, such as friendly grooming behavior or bite wounds, and is also spread from an infected mother cat to her kittens. Spaying eliminates this risk.. If your cat tested positive, please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

Feline Leukemia Virus (FeLV) is a viral infection in cats. There is no effective treatment to eliminate the infection. FeLV is spread from cat to cat through saliva, such as friendly grooming behavior or bite wounds, and is also spread from an infected mother cat to her kittens. Spaying eliminates this risk. If your cat tested positive, please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

Feline Immunodeficiency Virus (FIV) is a cats-only virus that is very similar to HIV in humans. FIV infection cannot be eliminated in a cat, but the cat may live symptom-free for a very long time or a normal life span. This virus is most commonly spread through bite wounds. Spaying or neutering greatly decreases fighting, and therefore helps to decrease risk of FIV infection to other cats. Please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

Feline Immunodeficiency Virus (FIV) is a cats-only virus that is very similar to HIV in humans. FIV infection cannot be eliminated in a cat, but the cat may live symptom-free for a very long time or a normal life span. This virus is most commonly spread through bite wounds. Spaying or neutering greatly decreases fighting, and therefore helps to decrease risk of FIV infection to other cats. Please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

Feline Immunodeficiency Virus (FIV) is a cats-only virus that is very similar to HIV in humans. FIV infection cannot be eliminated in a cat, but the cat may live symptom-free for a very long time or a normal life span. This virus is most commonly spread through bite wounds. Spaying or neutering greatly decreases fighting, and therefore helps to decrease risk of FIV infection to other cats. Please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

Feline Immunodeficiency Virus (FIV) is a cats-only virus that is very similar to HIV in humans. FIV infection cannot be eliminated in a cat, but the cat may live symptom-free for a very long time or a normal life span. This virus is most commonly spread through bite wounds. Spaying or neutering greatly decreases fighting, and therefore helps to decrease risk of FIV infection to other cats. Please see your regular veterinarian for confirmation of positive test status and health care recommendations.

---

- Your dog has been prescribed Gabapentin to reduce pain & anxiety, & provide mild sedation.
  - The sedation can help restrict activity to aid in healing after surgery.
  - Please follow all label directions for administering medication.
  - If your pet is too sedate or has any vomiting or diarrhea on this medication, stop giving the medication & contact us at **855-4FIX-AVL**. This helpline is open 24 hours.
- 

- Your dog has been prescribed Gabapentin to reduce pain & anxiety, & provide mild sedation.
  - The sedation can help restrict activity to aid in healing after surgery.
  - Please follow all label directions for administering medication.
  - If your pet is too sedate or has any vomiting or diarrhea on this medication, stop giving the medication & contact us at **855-4FIX-AVL**. This helpline is open 24 hours.
- 

- Your dog has been prescribed Gabapentin to reduce pain & anxiety, & provide mild sedation.
  - The sedation can help restrict activity to aid in healing after surgery.
  - Please follow all label directions for administering medication.
  - If your pet is too sedate or has any vomiting or diarrhea on this medication, stop giving the medication & contact us at **855-4FIX-AVL**. This helpline is open 24 hours.
- 

- Your dog has been prescribed Gabapentin to reduce pain & anxiety, & provide mild sedation.
- The sedation can help restrict activity to aid in healing after surgery.
  - Please follow all label directions for administering medication.
  - If your pet is too sedate or has any vomiting or diarrhea on this medication, stop giving the medication & contact us at **855-4FIX-AVL**. This helpline is open 24 hours.
-

Heartworm disease is an infection acquired from mosquitos that results in worms living in a dog or cat's heart, lungs, and surrounding blood vessels. A monthly pill easily prevents heartworm infection; once acquired, the infection can also, more expensively, be treated. Without treatment, this disease will slowly progress until it becomes fatal. Please see your regular veterinarian for treatment recommendations.

---

Heartworm disease is an infection acquired from mosquitos that results in worms living in a dog or cat's heart, lungs, and surrounding blood vessels. A monthly pill easily prevents heartworm infection; once acquired, the infection can also, more expensively, be treated. Without treatment, this disease will slowly progress until it becomes fatal. Please see your regular veterinarian for treatment recommendations.

---

Heartworm disease is an infection acquired from mosquitos that results in worms living in a dog or cat's heart, lungs, and surrounding blood vessels. A monthly pill easily prevents heartworm infection; once acquired, the infection can also, more expensively, be treated. Without treatment, this disease will slowly progress until it becomes fatal. Please see your regular veterinarian for treatment recommendations.

---

Heartworm disease is an infection acquired from mosquitos that results in worms living in a dog or cat's heart, lungs, and surrounding blood vessels. A monthly pill easily prevents heartworm infection; once acquired, the infection can also, more expensively, be treated. Without treatment, this disease will slowly progress until it becomes fatal. Please see your regular veterinarian for treatment recommendations.

---



- Please apply an ice pack, wrapped in a clean cloth, to the incision and surrounding area 2 to 3 times a day.
  - Apply for 10 minutes at a time to alleviate swelling.
  - Avoid placing ice pack directly on skin.
  - Discontinue icing 48 hours after surgery.
- 

- Please apply an ice pack, wrapped in a clean cloth, to the incision and surrounding area 2 to 3 times a day.
  - Apply for 10 minutes at a time to alleviate swelling.
  - Avoid placing ice pack directly on skin.
  - Discontinue icing 48 hours after surgery.
- 

- Please apply an ice pack, wrapped in a clean cloth, to the incision and surrounding area 2 to 3 times a day.
  - Apply for 10 minutes at a time to alleviate swelling.
  - Avoid placing ice pack directly on skin.
  - Discontinue icing 48 hours after surgery.
- 

- Please apply an ice pack, wrapped in a clean cloth, to the incision and surrounding area 2 to 3 times a day.
  - Apply for 10 minutes at a time to alleviate swelling.
  - Avoid placing ice pack directly on skin.
  - Discontinue icing 48 hours after surgery.
-

- Your dog has been prescribed Meloxicam, which is a non-steroidal anti-inflammatory drug (NSAID). It will provide pain control and help to decrease swelling.
- Please follow all label directions, and give the medication with a meal.
- Do not give this medication if your dog is already on an NSAID or any form of steroid.
- Possible side effects include loss of appetite, vomiting, diarrhea, dark stools, and/or decreased activity.
- If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.

Updated: 6/27/16

---

- Your dog has been prescribed Meloxicam, which is a non-steroidal anti-inflammatory drug (NSAID). It will provide pain control and help to decrease swelling.
- Please follow all label directions, and give the medication with a meal.
- Do not give this medication if your dog is already on an NSAID or any form of steroid.
- Possible side effects include loss of appetite, vomiting, diarrhea, dark stools, and/or decreased activity.
- If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.

Updated: 6/27/16

---

- Your dog has been prescribed Meloxicam, which is a non-steroidal anti-inflammatory drug (NSAID). It will provide pain control and help to decrease swelling.
- Please follow all label directions, and give the medication with a meal.
- Do not give this medication if your dog is already on an NSAID or any form of steroid.
- Possible side effects include loss of appetite, vomiting, diarrhea, dark stools, and/or decreased activity.
- If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.

Updated: 6/27/16

---

Your pet was found to be nauseous while staying with us overnight. Nausea is a common occurrence after surgery involving anesthesia, and usually resolves on its own. Please feed smaller amounts of your pet's normal food for a day or so until their system returns to normal. We have given your pet one or more of the following treatments to help with nausea and hydration:

- \_\_\_\_\_ mL Cerenia (anti-nausea)
- \_\_\_\_\_ mL Famotidine (anti-acid)
- \_\_\_\_\_ mL subcutaneous fluids (for hydration)

If vomiting continues longer than 12-24 hours, or if you have any questions, please call **(828) 252-2079** (local office hours) to schedule a recheck, or **(855) 434-9285** (after hours helpline).

---

Your pet was found to be nauseous while staying with us overnight. Nausea is a common occurrence after surgery involving anesthesia, and usually resolves on its own. Please feed smaller amounts of your pet's normal food for a day or so until their system returns to normal. We have given your pet one or more of the following treatments to help with nausea and hydration:

- \_\_\_\_\_ mL Cerenia (anti-nausea)
- \_\_\_\_\_ mL Famotidine (anti-acid)
- \_\_\_\_\_ mL subcutaneous fluids (for hydration)

If vomiting continues longer than 12-24 hours, or if you have any questions, please call **(828) 252-2079** (local office hours) to schedule a recheck, or **(855) 434-9285** (after hours helpline).

---

Your pet was found to be nauseous while staying with us overnight. Nausea is a common occurrence after surgery involving anesthesia, and usually resolves on its own. Please feed smaller amounts of your pet's normal food for a day or so until their system returns to normal. We have given your pet one or more of the following treatments to help with nausea and hydration:

- \_\_\_\_\_ mL Cerenia (anti-nausea)
- \_\_\_\_\_ mL Famotidine (anti-acid)
- \_\_\_\_\_ mL subcutaneous fluids (for hydration)

If vomiting continues longer than 12-24 hours, or if you have any questions, please call **(828) 252-2079** (local office hours) to schedule a recheck, or **(855) 434-9285** (after hours helpline).

Your pet should be on strict rest for \_\_\_\_\_ days.

No running, jumping, or rough playing. Only short, leashed walks should be provided, in order to urinate or defecate. We recommend keeping them confined to a small room or a kennel to ensure activity restriction. If using either a small room or kennel, your pet will initially need to be supervised to ensure their safety.

---

Your pet should be on strict rest for \_\_\_\_\_ days.

No running, jumping, or rough playing. Only short, leashed walks should be provided, in order to urinate or defecate. We recommend keeping them confined to a small room or a kennel to ensure activity restriction. If using either a small room or kennel, your pet will initially need to be supervised to ensure their safety.

---

Your pet should be on strict rest for \_\_\_\_\_ days.

No running, jumping, or rough playing. Only short, leashed walks should be provided, in order to urinate or defecate. We recommend keeping them confined to a small room or a kennel to ensure activity restriction. If using either a small room or kennel, your pet will initially need to be supervised to ensure their safety.

---

Your pet should be on strict rest for \_\_\_\_\_ days.

No running, jumping, or rough playing. Only short, leashed walks should be provided, in order to urinate or defecate. We recommend keeping them confined to a small room or a kennel to ensure activity restriction. If using either a small room or kennel, your pet will initially need to be supervised to ensure their safety.

A doctor has noted that your animal has an abnormal skin condition. Oftentimes, fleas are a major factor in this. Your animal may also have allergies, yeast, and/or bacterial infections. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable. Good over-the-counter flea products for dogs and cats include *Advantage* and *Frontline* (which will treat ticks as well).

---

A doctor has noted that your animal has an abnormal skin condition. Oftentimes, fleas are a major factor in this. Your animal may also have allergies, yeast, and/or bacterial infections. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable. Good over-the-counter flea products for dogs and cats include *Advantage* and *Frontline* (which will treat ticks as well).

---

A doctor has noted that your animal has an abnormal skin condition. Oftentimes, fleas are a major factor in this. Your animal may also have allergies, yeast, and/or bacterial infections. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable. Good over-the-counter flea products for dogs and cats include *Advantage* and *Frontline* (which will treat ticks as well).

---

A doctor has noted that your animal has an abnormal skin condition. Oftentimes, fleas are a major factor in this. Your animal may also have allergies, yeast, and/or bacterial infections. Please see your regular veterinarian so that your animal may be treated appropriately and become more comfortable. Good over-the-counter flea products for dogs and cats include *Advantage* and *Frontline* (which will treat ticks as well).

---

## Humane Alliance

A Program of the **ASPCA**

## Client Trazadone (Mild Sedative) Handout

- Your pet has been prescribed Trazadone, a mild sedative to help keep him/her calm. Please follow all label directions.
  - Do not give this medication if your pet is already on medication to reduce fear or anxiety.
  - Possible side effects include anxiousness, vomiting, diarrhea, seizures, hypertension and/or cardiac arrhythmia.
  - If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.
- 

## Humane Alliance

A Program of the **ASPCA**

## Client Trazadone (Mild Sedative) Handout

- Your pet has been prescribed Trazadone, a mild sedative to help keep him/her calm. Please follow all label directions.
  - Do not give this medication if your pet is already on medication to reduce fear or anxiety.
  - Possible side effects include anxiousness, vomiting, diarrhea, seizures, hypertension and/or cardiac arrhythmia.
  - If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.
- 

## Humane Alliance

A Program of the **ASPCA**

## Client Trazadone (Mild Sedative) Handout

- Your pet has been prescribed Trazadone, a mild sedative to help keep him/her calm. Please follow all label directions.
  - Do not give this medication if your pet is already on medication to reduce fear or anxiety.
  - Possible side effects include anxiousness, vomiting, diarrhea, seizures, hypertension and/or cardiac arrhythmia.
  - If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.
- 

## Humane Alliance

A Program of the **ASPCA**

## Client Trazadone (Mild Sedative) Handout

- Your pet has been prescribed Trazadone, a mild sedative to help keep him/her calm. Please follow all label directions.
- Do not give this medication if your pet is already on medication to reduce fear or anxiety.
- Possible side effects include anxiousness, vomiting, diarrhea, seizures, hypertension and/or cardiac arrhythmia.
- If you see any of these side effects, stop giving the medication and contact Humane Alliance at **855-4FIX-AVL**. This help line is open 24 hours.

- Run hot water over a wash-cloth.
  - Wring out wash-cloth so it is damp, but not dripping wet.
  - We want the wash-cloth to be as warm as possible, BUT if it is too hot for your face, it is too hot for your pet.
  - Hold the warm wash-cloth against the surgery area for \_\_\_\_\_ minutes, \_\_\_\_\_ times a day for \_\_\_ days.
  - The warm compress will help healing and make your pet more comfortable.
- 

- Run hot water over a wash-cloth.
  - Wring out wash-cloth so it is damp, but not dripping wet.
  - We want the wash-cloth to be as warm as possible, BUT if it is too hot for your face, it is too hot for your pet.
  - Hold the warm wash-cloth against the surgery area for \_\_\_\_\_ minutes, \_\_\_\_\_ times a day for \_\_\_ days.
  - The warm compress will help healing and make your pet more comfortable.
- 

- Run hot water over a wash-cloth.
  - Wring out wash-cloth so it is damp, but not dripping wet.
  - We want the wash-cloth to be as warm as possible, BUT if it is too hot for your face, it is too hot for your pet.
  - Hold the warm wash-cloth against the surgery area for \_\_\_\_\_ minutes, \_\_\_\_\_ times a day for \_\_\_ days.
  - The warm compress will help healing and make your pet more comfortable.
-